

Self-Duality of Super D3-brane Action on $AdS_5 \times S^5$ Background

Tadahiko Kimura^{1†}

¹ *Department of Physics, Faculty of Science, Chiba University, Chiba 263, Japan*

[†] E-mail: kimura@cuphd.nd.chiba-u.ac.jp

Abstract

We show that the super D3-brane action on $AdS_5 \times S^5$ background recently constructed by Metsaev and Tseytlin is exactly invariant under the combination of the electric-magnetic duality transformation of the worldvolume gauge field and the $SO(2)$ rotation of $N = 2$ spinor coordinates. The action is shown to satisfy the Gaillard-Zumino duality condition, which is a necessary and sufficient condition for an action to be self-dual. Our proof needs no gauge fixing for the κ -symmetry.

1 Introduction

Recently much attention is concentrated on the connection between string theory and M-theory on $AdS_{p+2} \times S^{D-p-2}$ and extended superconformal theories in $p+1$ spacetime dimensions [1, 2, 3]. $AdS_5 \times S^5$ describes a maximally supersymmetric vacuum (besides flat space) of type IIB supergravity and is the near horizon geometry of the D3-brane solution. $AdS_4 \times S^7$ and $AdS_7 \times S^4$ are maximally supersymmetric vacua of 11-dimensional supergravity and are the near horizon geometries of the M2- and M5-branes, respectively.

The actions of super p -branes in these near horizon background are described by a Dirac-Born-Infeld (DBI) type term and a Wess-Zumino term, and are invariant under local diffeomorphisms of the worldvolume, the κ -symmetry and the global isometries of the background.

Recently the super p -brane actions on $AdS_{p+2} \times S^{D-p-2}$ background geometries have been constructed for the superstring [4, 5, 6, 7, 8] and the D3-brane [9] on the $AdS_5 \times S^5$ background, for the M2-brane on the $AdS_4 \times S^7$ solution [10, 11] and M5-brane on the $AdS_7 \times S^4$ solution [12].

On the other hand electric-magnetic duality relations among various super p -brane effective actions on the flat background have been found in semiclassical approximation [13, 14] and for some cases on quantum level [15, 16, 17]. It will be of great interest to investigate whether these duality relations hold valid for super p -brane actions on AdS background. In some recent literatures the quantum equivalence of super D-string and F-string on $AdS_5 \times S^5$ and the semiclassical self-duality of super D3-brane on $AdS_5 \times S^5$ have already been discussed [18, 19]. However in these works proofs of the self-duality relied on a specific gauge choice for the κ -symmetry.

The purpose of the present paper is to show that the super D3-brane action on $AdS_5 \times S^5$ constructed by Metsaev and Tseytlin [9] is exactly self-dual. The action is shown to satisfy the Gaillard-Zumino(GZ) duality condition [20, 21, 22]. Igarashi et.al. have shown that the GZ-condition is actually necessary and sufficient condition for the theory to be self-dual [23] and they have shown that the super D3-brane action on the flat background indeed satisfies the GZ-duality condition [17].

This paper is organized as follows. In the next section a brief review of the super D3-brane action on $AdS_5 \times S^5$ constructed by Metsaev and Tseytlin by the coset formalism [4, 9] is given. In section 3 the super D3-brane action on $AdS_5 \times S^5$ is shown to satisfy the GZ duality condition, thereby establishing its exact self-duality without resort to any semiclassical approximation.

2 Super D3-brane Action on $AdS_5 \times S^5$

In this section, for later use, we briefly review the D3-brane action constructed by Metsaev and Tseytlin [9].

Supervielbein superfields on the $AdS_5 \times S^5$ background are explicitly constructed by the coset formalism [4, 5]. In the explicit coset parametrization $G(X^{\hat{m}}, \theta) = g(X^{\hat{m}})e^{\theta Q}$, where $X^M = (X^{\hat{m}}, \theta)$ and Q are $N = 2$ superspace coordinate and su-

percharge, respectively, the supervielbein $L^A = (L^{\hat{a}}, L)$ and superconnection $L^{\hat{a}\hat{b}}$ are explicitly given as follows;

$$L^{\hat{a}} = e^{\hat{a}} - 2i\bar{\theta}\Gamma^{\hat{a}}W(\theta)D\theta, L = V(\theta)D\theta, \quad (2.1)$$

$$L^{ab} = \omega^{ab} + 2\bar{\theta}i\tau^{(2)}\Gamma^{ab}\sigma_-W(\theta)D\theta, L^{a'b'} = \omega^{a'b'} + 2\bar{\theta}i\tau^{(2)}\Gamma^{a'b'}\sigma_-W(\theta)D\theta, \quad (2.2)$$

where a, b (a', b') are AdS_5 (S^5) coordinate indices and run 0,1,2,3,4 (5,6,7,8,9), and $D\theta, V(\theta)$ and $W(\theta)$ are defined by

$$D\theta = (d + \frac{1}{4}\omega^{\hat{a}\hat{b}}\Gamma^{\hat{a}\hat{b}} + \frac{1}{2}\tau^{(2)}e^{\hat{a}}\sigma_+\Gamma^{\hat{a}})\theta. \quad (2.3)$$

$$V(\theta) = \frac{\sinh \sqrt{m(\theta)}}{\sqrt{m(\theta)}} = 1 + \frac{1}{3!}m(\theta) + \frac{1}{5!}m(\theta)^2 + \dots, \quad (2.4)$$

$$W(\theta) = \frac{\cosh \sqrt{m(\theta)} - 1}{m(\theta)} = \frac{1}{2!} + \frac{1}{4!}m(\theta) + \frac{1}{6!}m(\theta)^2 + \dots, \quad (2.5)$$

$$m(\theta) = -\sigma_+\Gamma^{\hat{a}}i\tau^{(2)}\theta\bar{\theta}\Gamma^{\hat{a}} + \frac{1}{2}\Gamma^{ab}\theta\bar{\theta}i\tau^{(2)}\Gamma^{ab}\sigma_- - \frac{1}{2}\Gamma^{a'b'}\theta\bar{\theta}i\tau^{(2)}\Gamma^{a'b'}\sigma_-. \quad (2.6)$$

In the above equations Pauli matrices $\tau^{(i)}$ operate on internal $N = 2$ space and Pauli matrices σ_i stand for 32×32 matrices $1 \times 1 \times \sigma_i$.

The D3-brane action depends on the coset superspace coordinates $X^M = (X^{\hat{m}}, \theta)$ and U(1) world-volume gauge field strength $F = dA = \frac{1}{2}F_{ij}d\sigma^i \wedge d\sigma^j$, $F_{ij} = \partial_i A_j - \partial_j A_i$, where i and j run over the worldvolume indices 0,1,2,3. The reparametrization and κ -symmetry invariant super D3-brane action in the $AdS_5 \times S^5$ background consists of two terms; i.e. the Dirac-Born-Infeld(DBI) term and the Wess-Zumino term:

$$S = S_{DBI} + S_{WZ}. \quad (2.7)$$

The DBI term is given by

$$S_{DBI} = - \int_{M_4} d^4\sigma \sqrt{-\det(G_{ij} + \mathcal{F}_{ij})}, \quad (2.8)$$

where G_{ij} is the supersymmetric induced worldvolume metric defined by

$$G_{ij} = L^{\hat{a}}_i L^{\hat{a}}_j = \partial_i X^M L^{\hat{a}}_M \partial_j X^N L^{\hat{a}}_N, L^{\hat{a}} = d\sigma^i L^{\hat{a}}_i, \quad (2.9)$$

and two-form superfield $\mathcal{F}$ is the supersymmetric extension of the U(1) worldvolume gauge field strength defined by

$$\mathcal{F} = dA + \Omega^{(3)}, \quad (2.10)$$

where two-form $\Omega^{(i)}$ is defined by

$$\Omega^{(i)} = 2i \int_0^1 ds \bar{\theta} \hat{L}_s \tau^{(i)} L_s, \quad (2.11)$$

and $L_s^{\hat{a}} \equiv L^{\hat{a}}(x, s\theta)$, $L_s \equiv L(x, s\theta)$ and $\hat{L}_s \equiv L_s^{\hat{a}} \Gamma^{\hat{a}}$.

The Wess-Zumino term is given by

$$S_{WZ} = \int_{M_4} \Omega_4 = \int_{M_5} H_5, \quad (2.12)$$

here

$$\begin{aligned} H_5 &= d\Omega_4 \\ &= i\bar{L} \wedge \left(\frac{1}{6} \hat{L} \wedge \hat{L} \wedge \hat{L} i\tau^{(2)} + \mathcal{F} \wedge \hat{L} \tau^{(1)} \right) \wedge L \\ &+ \frac{1}{30} (\epsilon^{a_1 \dots a_5} L^{a_1} \wedge \dots \wedge L^{a_5} + \epsilon^{a'_1 \dots a'_5} L^{a'_1} \wedge \dots \wedge L^{a'_5}). \end{aligned} \quad (2.13)$$

It is convenient to rewrite Ω_4 in the following form;

$$\Omega_4 = C_4 + \Omega^{(1)} \wedge \mathcal{F}, \quad (2.14)$$

then we obtain a differential equation for C_4

$$\begin{aligned} dC_4 + \Omega^{(1)} \wedge d\Omega^{(3)} &= i\frac{1}{6} \bar{L} \wedge \hat{L} \wedge \hat{L} \wedge \hat{L} i\tau^{(2)} \wedge L \\ &+ \frac{1}{30} (\epsilon^{a_1 \dots a_5} L^{a_1} \wedge \dots \wedge L^{a_5} + \epsilon^{a'_1 \dots a'_5} L^{a'_1} \wedge \dots \wedge L^{a'_5}). \end{aligned} \quad (2.15)$$

The equation (2.13) or (2.15) can be explicitly integrated by using the Maurer-Cartan equations and the $\theta \rightarrow s\theta$ trick and the result is given in [9]. However for our purpose the explicit form of C_4 is not necessary.

As it can be seen from (2.1)-(2.6), the super D3-brane action on $AdS_5 \times S^5$ is a complicated function of supercoordinate θ . However to prove the self-duality of the action (2.7) we need no gauge fixing condition to simplify the fermionic dependence of the supervielbeins and superconnections.

3 Self-Duality of Super D3-brane Action on $AdS_5 \times S^5$

In this section we show that the super D3-brane action on $AdS_5 \times S^5$ reviewed in the last section satisfies the Gaillard and Zumino duality-condition, thereby establishing its exact self-duality without resort to any semiclassical approximation.

First let us recall what is the GZ duality condition. Given a generic Lagrangian density $\mathcal{L}(F_{\mu\nu}, g_{\mu\nu}, \phi^A) = \sqrt{-g} L(F_{\mu\nu}, g_{\mu\nu}, \phi^A)$ in four dimensional spacetime which

contains a $U(1)$ gauge field strength $F_{\mu\nu}$, gravitational field $g_{\mu\nu}$ and generic matter fields Φ^A , the constructive relation is given by

$$\tilde{K}_{\mu\nu} \equiv \frac{\partial L}{\partial F_{\mu\nu}}, \quad (3.1)$$

where the Hodge dual components for the anti-symmetric tensor $K_{\mu\nu}$ are defined by

$$\tilde{K}_{\mu\nu} \equiv \frac{1}{2}\eta_{\mu\nu}^{\rho\sigma}K_{\rho\sigma}, \quad \tilde{\tilde{K}}_{\mu\nu} = -K_{\mu\nu}, \quad (3.2)$$

where $\eta_{\mu\nu\rho\sigma} = \sqrt{-g}\epsilon_{\mu\nu\rho\sigma}$, $\epsilon^{0123} = 1$ and the signature of $g_{\mu\nu}$ is $(-, +, +, +)$.

If one defines the infinitesimal $SO(2)$ duality transformation by

$$\begin{aligned} \delta F_{\mu\nu} &= \lambda K_{\mu\nu}, \quad \delta K_{\mu\nu} = -\lambda F_{\mu\nu}, \\ \delta \Phi^A &= \xi^A(\Phi), \end{aligned} \quad (3.3)$$

then the consistency of the constructive relation (3.1) and the invariance of the field equations under this $SO(2)$ duality transformation require the following condition:

$$\frac{\lambda}{4}(F_{\mu\nu}\tilde{F}^{\mu\nu} + K_{\mu\nu}\tilde{K}^{\mu\nu}) + \delta_\Phi \mathcal{L} = 0, \quad (3.4)$$

and the invariance of the energy-momentum tensor requires $\delta g_{\mu\nu} = 0$. We call the condition (3.4) the GZ self-duality condition [20, 21].

It has been known that the $SO(2)$ duality is lifted to the $SL(2, R)$ duality by introducing a dilaton ϕ and an axion χ [24].

Moreover it has also been shown that the GZ condition (3.4) is actually the necessary and sufficient condition for the action to be invariant under the duality transformation (3.3) [23]. Therefore if one can show an action to satisfy the condition (3.4) under the transformation (3.3) with suitable transformation rule for matter fields, then one establishes the exact self-duality of the theory described by this action without resort to any semiclassical approximation.

Now let us show that the super D3-brane action on $AdS_5 \times S^5$ reviewed in the preceding section satisfies the GZ duality condition under the following $SO(2)$ duality transformation:

$$\begin{aligned} \delta F_{ij} &= \lambda K_{ij}, \quad \delta K_{ij} = -\lambda F_{ij}, \\ \delta \theta &= -\lambda \frac{i\tau^{(2)}}{2}\theta, \quad \delta \bar{\theta} = \lambda \bar{\theta} \frac{i\tau^{(2)}}{2}, \quad \delta X = 0. \end{aligned} \quad (3.5)$$

The $N = 2$ spinor coordinates transform as an $SO(2)$ doublet.

Let us first examine how various quantities in the action (2.7) transform under the above $SO(2)$ duality rotation. The matrix $m(\theta)$ defined in (2.6) transforms as an adjoint representation of $SO(2)$, and therefore matrices $V(\theta)$ and $W(\theta)$ obey the same transformation properties as $m(\theta)$;

$$\delta m(\theta) = -\lambda \left[\frac{i}{2}\tau^{(2)}, m(\tau) \right], \quad \delta V(\theta) = -\lambda \left[\frac{i}{2}\tau^{(2)}, V(\tau) \right], \quad \delta W(\theta) = -\lambda \left[\frac{i}{2}\tau^{(2)}, W(\tau) \right]. \quad (3.6)$$

From these transformation properties, we can easily show that the supervielbeins transform as follows;

$$\delta L^{\hat{a}} = 0, \delta L = -\lambda \frac{i\tau^{(2)}}{2} L, \delta \bar{L} = \lambda \bar{L} \frac{i\tau^{(2)}}{2}, \quad (3.7)$$

namely the $N = 2$ spinor components of the supervielbeins L and $\bar{L}$ transform as $SO(2)$ doublets.

From (3.5) and (3.7), we find $\Omega^{(1,3)} = 2i \int_0^1 ds \bar{\theta} \hat{L}_s \tau^{(1,3)} L_s$ defined in (2.11) transform as an $SO(2)$ doublet and $\Omega^{(2)}$ is an $SO(2)$ singlet:

$$\delta \Omega^{(1)} = \lambda \Omega^{(3)}, \delta \Omega^{(3)} = -\lambda \Omega^{(1)}, \delta \Omega^{(2)} = 0. \quad (3.8)$$

From (3.8) and (2.15) the transformation of C_4 is easily obtained. Noticing the right hand-side of (2.15) is invariant under $SO(2)$ rotation, the $SO(2)$ transformation of (2.15) yields

$$d\delta C_4 = -\frac{\lambda}{2} d(\Omega^{(1)} \wedge \Omega^{(1)} + \Omega^{(3)} \wedge \Omega^{(3)}). \quad (3.9)$$

Then we obtain the transformation rule for C_4 up to exact forms;

$$\delta C_4 = -\frac{\lambda}{2} (\Omega^{(1)} \wedge \Omega^{(1)} + \Omega^{(3)} \wedge \Omega^{(3)}) \quad (3.10)$$

Now we are ready to prove the duality symmetry of the super D3-brane action on $AdS_5 \times S^5$. First let us calculate $\frac{\lambda}{4}(F_{ij}\tilde{F}^{ij} + K_{ij}\tilde{K}^{ij})$. From the definition (3.1) and the action (2.7) we obtain

$$\begin{aligned} \tilde{K}^{ij} &= \frac{\partial L}{\partial F_{ij}} \\ &= \frac{\sqrt{-\det G_{ij}}}{\sqrt{-\det(G_{ij} + \mathcal{F}_{ij})}} (-\mathcal{F}^{ij} + \mathcal{T} \tilde{\mathcal{F}}^{ij}) + \tilde{\Omega}^{(1)ij}, \end{aligned} \quad (3.11)$$

where we have used the determinant formula for the four-by-four matrix:

$$\det(G_{ij} + \mathcal{F}_{ij}) = \det G_{ij} (1 + \frac{1}{2} \mathcal{F}_{ij} \mathcal{F}^{ij} - \mathcal{T}^2), \mathcal{T} \equiv \frac{1}{4} \mathcal{F}_{ij} \tilde{\mathcal{F}}^{ij}. \quad (3.12)$$

Taking the Hodge dual of (3.9), we find

$$K_{ij} = -\frac{1}{2} \eta_{ijkl} \tilde{K}^{kl} = \frac{\sqrt{-\det G_{ij}}}{\sqrt{-\det(G_{ij} + \mathcal{F}_{ij})}} (\tilde{\mathcal{F}}_{ij} + \mathcal{T} \mathcal{F}_{ij}) + \Omega_{ij}^{(1)}. \quad (3.13)$$

Then we obtain

$$\begin{aligned} \frac{\lambda}{4} (F_{ij} \tilde{F}^{ij} + K_{ij} \tilde{K}^{ij}) &= \frac{\lambda}{4} (-2\Omega_{ij}^{(3)} \tilde{F}^{ij} + 2\Omega_{ij}^{(1)} \tilde{K}^{ij} - \Omega_{ij}^{(1)} \tilde{\Omega}^{(1)ij} - \Omega_{ij}^{(3)} \tilde{\Omega}^{(3)ij}) \\ &= \frac{\lambda}{4} (-4\Omega^{(3)} F + 4\Omega^{(1)} K - 2\Omega^{(1)} \Omega^{(1)} - 2\Omega^{(3)} \Omega^{(3)}) \end{aligned} \quad (3.14)$$

In the above and below we omit $\wedge$ -symbol in products of forms.

Next let us calculate $\delta_\theta L$. In the language of differential forms,

$$\begin{aligned}
\delta_\theta L &= \frac{\partial L}{\partial \mathcal{F}} \delta \Omega^{(3)} + \mathcal{F} \delta \Omega^{(1)} + \delta C_4 \\
&= \lambda [-K \Omega^{(1)} + (F + \Omega^{(3)}) \Omega^{(3)} + \frac{1}{2} (\Omega^{(1)} \Omega^{(1)} - \Omega^{(3)} \Omega^{(3)})] \\
&= \lambda [-K \Omega^{(1)} + F \Omega^{(3)} + \frac{1}{2} (\Omega^{(1)} \Omega^{(1)} + \Omega^{(3)} \Omega^{(3)})].
\end{aligned} \tag{3.15}$$

It is clearly seen that the right-hand sides of (3-14) and (3-15) exactly cancel with each other and the GZ-duality condition is indeed satisfied. As we have proved the invariance of the action under the infinitesimal $SO(2)$ duality transformation, the action is also invariant under the finite $SO(2)$ duality transformation.

Before closing this section let us mention the extension from the $SO(2)$ duality to the $SL(2, R)$ duality by introducing a constant dilaton ϕ and an axion χ fields. We follow the same line of arguments given in [17]. According to the general method [24], let us define a new Lagrangian $\hat{L}(G, F, \theta, \phi, \chi)$ from the D3-brane Lagrangian $L(G, F, \theta)$ which obey the $SO(2)$ duality ;

$$\hat{L}(G, F, \theta, \phi, \chi) = L(G, e^{-\phi/2} F, \theta) + \frac{1}{4} \chi F \tilde{F}. \tag{3.16}$$

Then if one define $\hat{F} = e^{-\phi/2} F$ and $\hat{K}$ by taking the dual of $\frac{\partial L(G, \hat{F}, \theta)}{\partial \hat{F}}$, the background dependence is absorbed in the rescaled variables $(\hat{K}, \hat{F})$. These are related with the background dependent (K, F) by

$$\begin{pmatrix} K \\ F \end{pmatrix} = V \begin{pmatrix} \hat{K} \\ \hat{F} \end{pmatrix}, \tag{3.17}$$

$$V = e^{\frac{\phi}{2}} \begin{pmatrix} e^{-\phi} & \chi \\ 0 & 1 \end{pmatrix}. \tag{3.18}$$

Here V is a non-linear realization of $SL(2, R)/SO(2)$ transforming as

$$V \longrightarrow V' = \Lambda V O(\Lambda)^{-1} \tag{3.19}$$

Here Λ is a global $SL(2, R)$ matrix

$$\Lambda = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, R), ad - bc = 1, \tag{3.20}$$

and $O(\Lambda)$ is an $SO(2)$ transformation

$$O(\Lambda)^{-1} = \begin{pmatrix} \cos \lambda & \sin \lambda \\ -\sin \lambda & \cos \lambda \end{pmatrix} \in SO(2). \tag{3.21}$$

The condition that the form of V (3.17) is unchanged under the transformation (3.18) determines the $SO(2)$ rotation angle λ and the transformation rule of the background fields ϕ and χ ;

$$\tan \lambda = \frac{ce^{-\phi}}{c\chi + d}, \quad (3.22)$$

and

$$\tau \rightarrow \tau' = \frac{a\tau + b}{c\tau + d}; \tau \equiv \chi + ie^{-\phi}. \quad (3.23)$$

These results show that if the original Lagrangian $L(G, F, \theta)$ is invariant under the $SO(2)$ duality transformation the extended Lagrangian $\hat{L}(G, F, \theta, \phi, \chi)$ with a dilaton and an axion fields is invariant under the $SL(2, R)$ duality transformation of (K, F) and $\tau \equiv \chi + ie^{-\phi}$ and $SO(2)$ rotation of $N = 2$ spinor with rotation angle λ given by (3.21).

4 Discussion

We have shown that the super D3-brane action on $AdS_5 \times S^5$ background is exactly self-dual, by proving that the action satisfies the Gaillard and Zumino duality condition which is a necessary and sufficient condition for the action to be self-dual. It should be stressed that our proof of the self-duality needs not any gauge fixing choice for the κ -symmetry.

Although the super D3-brane action constructed by Metsaev and Tseytlin for which we have proved the self-duality is based on the explicit coset parametrization $G(X^{\hat{m}}, \theta) = g(X)e^{\theta Q}$, since the physical properties such as duality relations among super p-branes should not depend on the choice of the parametrization of manifolds, our result should be valid for any other parametrizations of coset manifold. In a recent paper [25] super p-brane theories based on a different parametrizations of the coset manifolds are proposed. It would be interesting to confirm that the super D3-brane action constructed based on this new coset parametrization is indeed self-dual.

Now the self-duality of the super D3-brane action has been proved for the flat and AdS background. Therefore we expect that the self-duality of super D3-brane and various duality relations among super p-branes will be proved in generic background geometries.

There are several effective actions of super D3-brane which are proposed to be manifestly duality symmetric [26, 27]. It would be interesting to investigate possible relations with the duality symmetry discussed in this paper.

We may think about various p -branes on $AdS_5 \times S^5$ background just as in the flat background. It would be interesting to construct various super p-brane actions on $AdS_5 \times S^5$ and investigate duality relations among them known on the flat background in semiclassical approximation [14].

It has been proposed that D-brane dualities are understandable as canonical transformation [28, 15, 17]. It would be also interesting to formulate dualities among super

D-branes and M-branes both on flat and AdS backgrounds as canonical transformations.

Acknowledgments

I would like to thank Y. Igarashi, K. Kamimura and I. Oda for explaining their works. This work is supported in part by the Grant-in-Aid for Research from the Ministry of Education, Science and Culture.

References

- [1] J.Maldacena, The large N limit of superconformal field theories and supergravity, hep-th/9711200.
- [2] S.S. Gupser, I.R. Klebanov and A.M. Polyakov, Phys. Lett. B428 (1998) 105; hep-th/9802109.
- [3] E. Witten, Anti de Sitter space and holography, hep-th/9802150.
- [4] R.R. Metsaev and A.A. Tseytlin, Type IIB Superstring Action in $AdS_5 \times S^5$ Background, hep-th/9805028.
- [5] R.Kallosh, J. Rahmfeld and A. Rajaraman, Near Horizon Superspace, hep-th/9805217.
- [6] R. Kallosh and Rahmfeld, The GS String action on $AdS_5 \times S^5$, hep-th/9808038.
- [7] R. Pesand, A κ Gauge Fixed Type IIB Superstring Action on $AdS_5 \times S^5$, hep-th/9808020.
- [8] R. Kallosh and A.A. Tsytlín, Simplifying supersymmetric action on $AdS_5 \times S^5$, hep-th/9808088.
- [9] R.R. Metsaev and A.A. Tseytlin, Supersymmetric D3 brane action in $AdS_5 \times S^5$, hep-th/9806095.
- [10] G. Dall'Agata, D. Fabbri, C. Fraser, P. Fré, P. Termonia and M. Trigiante, The $OSp(8|4)$ singleton action from the supermembrane, hep-th/9807115.
- [11] B. de Wit, K. Peeters, J. Plefka and A. Sevrin, The M-theory Two-Brane in $AdS_4 \times S^7$ and $AdS_7 \times S^4$, hep-th/9808052.
- [12] P. Claus, Super M-Brane action in $AdS_4 \times S^7$ and $AdS_7 \times S^4$, hep-th/9809045.
- [13] A. A. Tseytlin, Self-duality of Born-Infeld action and Dirichlet 3-brane of type IIB superstring theory, Nucl. Phys. B469 (1996) 51, hep-th/9602064.
- [14] M. Aganagic, J. Park, C. Popescu and J. H. Schwarz, Dual D-Brane Action, Nucl. Phys. B496 (1997) 215, hep-th/9702133.

- [15] Y. Igarashi, K. Itoh, K. Kamimura and R. Kuriki, Canonical Equivalence between Super D-string and Type II Superstring, JHTP 03 (1998) 002, hep-th/9801118.
- [16] I. Oda, Path Integral Equivalence between Super D-string and IIB Superstring, Phys. Lett. B430 (1998) 242, hep-th/9802152.
- [17] Y. Igarashi, K. Itoh and K. Kamimura, Self-duality in Super D3-brane Action, hep-th/9806161.
- [18] I. Oda, Super D -string Action on $AdS_5 \times S^5$, hep-th/9809076
- [19] I. Oda, $SL(2, Z)$ Self-duality of Super D3-brane Action on $AdS_5 \times S^5$, hep-th/9810024.
- [20] M.K. Gaillard and B. Zumino, Duality Rotations for Interaction Fields, Nucl. Phys. B193 (1981) 221-244.
- [21] M.K. Gaillard and B. Zumino, Self-Duality in Nonlinear Electromagnetism, hep-th/9705226; Nonlinear Electromagnetic Self-Duality and Legendre Transformations, hep-th/9712103.
- [22] G. W. Gibbons and D.A. Rasheed, Electric-magnetic duality rotations in non-linear electrodynamics, Nucl. Phys. B454 (1995) 185-206, hep-th/9506035.
- [23] Y. Igarashi, K. Itoh and K. Kamimura, Electric-Magnetic Duality Rotations and Invariance of Actions, hep-th/9806160.
- [24] G. W. Gibbons and D.A. Rasheed, $SL(2, R)$ invariance of non-linear electrodynamics coupled to an axion and a dilaton, Phys. Lett. B365 (1996) 46, hep-th/9509141.
- [25] P. Pasti, D. Sorokin and M. Tonin, On gauged-fixed superbrane actions in AdS superbackground, hep-th/9809213.
- [26] I. Bengtsson, Manifestly Duality in Born–Infeld Theory, hep-th/9612174; D. Berman, $SL(2, Z)$ duality of Born–Infeld Theory from non-linear self-dual electrodynamics in 6 dimensions, Phys. Lett. B409 (1997) 153, hep-th/9706208; M5 on a torus and the three brane, hep-th/9804115; A. Nurmagambetov, Duality-Symmetric Three–Brane and its Coupling to Type IIB Supergravity, hep-th/980157.
- [27] M.Cederwall and A. Westerberg, World-volume fields, $SL(2, Z)$ and duality:the type IIB 3-branes, J. High. Energy Phys. 02 (1998) 004, hep-th/9710007; P. K. Townsend, Membrane tension and manifestly IIB duality, Phys.Lett. B409 (1997) 131, hep-th/9705160; P. K. Townsend and M. Cederwall, The manifestly $SL(2, Z)$ covariant superstring, J. High Energy Phys. 09.03 (1997), hep-th/9709002.
- [28] Y. Lozano, D-Brane Dualities As Canonical Transformation, Phys. Lett. B399 (1997) 233, hep-th/9701186.